

STATUS AND PROSPECTS FOR LORENTZ AND CPT VIOLATION TESTS AT KLOE AND KLOE-2

ANTONIO DE SANTIS

*Dipartimento di Fisica, Università di Roma 'La Sapienza'
and I.N.F.N. Sezione di Roma*

P.le A. Moro, 2

I-00185 Rome, Italy

E-mail: antonio.desantis@roma1.infn.it

on behalf of the KLOE and KLOE-2 collaborations

The neutral kaon system offers a unique possibility to perform fundamental tests of CPT invariance. In this contribution the KLOE prospects for the measurements of CPT violation in the context of the Standard-Model Extension are presented together with a full description of the analysis method needed and with the perspective given by the KLOE-2 data-taking campaign.

1. The KLOE experiment

The KLOE experiment operates at DAΦNE, the Frascati ϕ -factory. DAΦNE is an e^+e^- collider running at a center of mass energy of ~ 1020 MeV, the mass of the ϕ meson. Positron and electron beams collide at an angle of π -25 mrad, producing ϕ mesons with small boost in the orbit plane ($p_x(\phi) \sim -15$ MeV).

The KLOE detector consists of a large cylindrical drift chamber (DC) surrounded by a lead-scintillating fiber electromagnetic calorimeter (EMC). A superconducting coil around the EMC provides a 0.52 T axial field. The DC¹ is 4 m in diameter and 3.3 m long and has 12,582 all-stereo tungsten sense wires. The chamber shell is made of carbon fiber-epoxy composite and the gas used is a 90% helium, 10% isobutane mixture. These features maximize transparency to photons and reduce $K_L \rightarrow K_S$ regeneration and multiple scattering. The position resolutions are $\sigma_{r,\phi} \sim 150 \mu\text{m}$ and $\sigma_z \sim 2$ mm. The momentum resolution is $\sigma(p_\perp)/p_\perp \sim 0.4\%$. The calorimeter² is divided into a barrel and two endcaps, and covers 98% of the solid angle. The modules are read-out at both ends by photomultipliers, both in am-

plitude and time for a total of 2440 cells per side arranged in five layers. Cells close in time and space are grouped into calorimeter clusters. The cluster energy E is the sum of the cell energies. The cluster time T and position $\vec{R}$ are energy-weighted averages. Energy and time resolutions are $\sigma_E/E = 5.7\%/\sqrt{E(\text{GeV})}$ and $\sigma_t = 57\text{ps}/\sqrt{E(\text{GeV})} \oplus 100\text{ ps}$, respectively.

Actually KLOE has already acquired 2.5 fb^{-1} of data and a new extensive campaign of data taking is starting aiming at an integrated luminosity of 25 fb^{-1} .

The cross section for the process $e^+e^- \rightarrow \phi$ production is $3.3\text{ }\mu\text{b}$ and the ϕ meson decays into $K_0\bar{K}_0$ with a branching fraction of $\sim 34\%$. The initial state of the kaon pair is produced via strong interaction with quantum numbers $J^{PC} = 1^{--}$:

$$|i\rangle = \frac{|K_0\rangle|\bar{K}_0\rangle - |\bar{K}_0\rangle|K_0\rangle}{\sqrt{2}} = \mathcal{N}(|K_S\rangle|K_L\rangle - |K_S\rangle|K_L\rangle), \quad (1)$$

where $|K_S/K_L\rangle = (1 + \varepsilon_{S/L})|K_0\rangle \pm (1 - \varepsilon_{S/L})|\bar{K}_0\rangle$ and $\varepsilon_{S/L} = \varepsilon \pm \delta$. The two kaons are in an antisymmetric correlated state and the time evolution of their state in two different final states f_1 and f_2 for the two kaons as a function of the difference of decay time ($\Delta t = t_2 - t_1$) can be expressed as the decay intensity as a function of Δt :

$$I_{f_1 f_2}(\Delta t) \propto |\eta_1|e^{-\Gamma_L \Delta t} + |\eta_2|e^{-\Gamma_S \Delta t} - 2e^{-\frac{(\Gamma_S + \Gamma_L)}{2} \Delta t} \cos(\Delta m \Delta t + \phi_2 - \phi_1), \quad (2)$$

where

$$\eta_j = \langle f_j | K_L \rangle / \langle f_j | K_S \rangle = |\eta_j| e^{i\phi_j}. \quad (3)$$

2. CPT and Lorentz symmetry breaking

A general theoretical possibility for CPT violation is based on spontaneous breaking of Lorentz symmetry, as developed by Kostelecký:³⁻⁵ the Standard-Model Extension (SME).

In SME for neutral kaons, CPT manifests to lowest order only in the CPT violation parameter δ , and exhibits a dependence on the 4-momentum of the kaon:

$$\delta \approx i \sin \phi_{SW} e^{i\phi_{SW}} \gamma_K (\Delta a_0 - \vec{\beta}_K \cdot \Delta \vec{a}) / \Delta m, \quad (4)$$

where γ_K and $\vec{\beta}_K$ are the kaon boost factor and velocity in the observer frame, and Δa_μ are the CPT violation coefficients for the two valence quarks in the kaon.

The time dependence arising from the rotation of the Earth can be explicitly displayed in Eq. (4) by choosing a three-dimensional basis $(\hat{X}, \hat{Y}, \hat{Z})$ in a nonrotating frame, with the $\hat{Z}$ axis along the Earth's rotation axis, and a basis $(\hat{x}, \hat{y}, \hat{z})$ for the laboratory frame.⁴ The CPT violating parameter δ may then be expressed as:

$$\delta(\vec{p}_K, t) = \frac{i \sin \phi_{SW} e^{i\phi_{SW}}}{\Delta m} \gamma_K \left[\Delta a_0 + \beta_K \Delta a_Z (\cos \vartheta \cos \chi - \sin \vartheta \cos \phi \sin \chi) \right. \\ \left. - \beta_K \Delta a_X \sin \vartheta \sin \phi \sin \Omega t \right. \\ \left. + \beta_K \Delta a_X (\cos \vartheta \sin \chi + \sin \vartheta \cos \phi \sin \chi) \cos \Omega t \right. \\ \left. + \beta_K \Delta a_Y (\cos \vartheta \sin \chi + \sin \vartheta \cos \phi \sin \chi) \sin \Omega t \right. \\ \left. - \beta_K \Delta a_Y \sin \vartheta \sin \phi \cos \Omega t \right], \quad (5)$$

where t is the sidereal time, Ω is the Earth's sidereal frequency and χ is the angle between the Earth rotation axis and the $\hat{z}$ direction in the laboratory frame.

Since inside the KLOE detector, kaons are produced in almost all direction, it is possible to use the experiment like a telescope to explore any direction in the space by using two different method to measure the Δa_μ parameters of Eq. (4):

- by studying the sidereal time variation of the semileptonic asymmetry of the K_L and K_S . A preliminary measurement of the Δa_0 has been already performed with KLOE data using the difference between the semileptonic asymmetries of K_S and K_L : $\Delta a_0 = (0.4 \pm 1.8) \times 10^{-17} \text{ GeV}$.⁶
- by studying the quantum interferometry between the two kaons, Eq. (2). In this case, two different final state can be considered: $(\pi l^+ \nu, \pi l^- \bar{\nu})$ and $(\pi^+ \pi^-, \pi^+ \pi^-)$.

In the following we will focus on interferometry with the channel $\phi \rightarrow K_S K_L \rightarrow \pi^+ \pi^- \pi^+ \pi^-$. Detailed discussion concerning other methods can be found elsewhere.⁶ Using this channel the two kaons decay both in the same final state, but have opposite momenta and experience different direction in the space. For instance, we can distinguish between the two kaons by their forward or backward emission. In this way it is possible to define as K_+ (K_-) the one going in the forward (backward) direction with $\cos(\vartheta) > (<) 0$. The ratio of amplitudes in Eq. (3) is $\eta_\pm = \varepsilon - \delta(\vec{p})$. Since the kaon momenta are opposite, a small asymmetry in the decay intensity

$I_{\pm}(\Delta t)$ will appear and the following ratio will be different from zero:

$$A(\Delta t) = \frac{I_{\pm}(\Delta t > 0) - I_{\pm}(\Delta t < 0)}{I_{\pm}(\Delta t > 0) + I_{\pm}(\Delta t < 0)}. \quad (6)$$

The above asymmetry for $\Delta t \gg \tau_S$ tends to zero, because ε and δ are 90° out of phase:⁷

$$A(\Delta t \tau_S) \simeq -2\Re\left(\frac{\delta}{\varepsilon}\right) \sim 0, \quad (7)$$

while for $|\Delta t| \leq 5\tau_S$

$$A(|\Delta t| 5\tau_S) \propto -2\Im m\left(\frac{\delta}{\varepsilon}\right) \quad (8)$$

and therefore to $\Delta a_{X,Y,Z}$.

With analysis based on 1 fb^{-1} we get a preliminary result on $\Delta a_{X,Y,Z}$:⁸

$$\begin{aligned} \Delta a_X &= (-6.3 \pm 6.0) \times 10^{-18} \text{ GeV}, \\ \Delta a_Y &= (2.8 \pm 5.9) \times 10^{-18} \text{ GeV}, \\ \Delta a_Z &= (2.4 \pm 9.7) \times 10^{-18} \text{ GeV}. \end{aligned} \quad (9)$$

This analysis scheme using the simple forward backward asymmetry is not sensitive to Δa_0 and its effect is washed out in the asymmetry Eq. (5) and Eq. (6). As shown in Eq. (4) this parameter is coupled only with the γ_K factor. Since at DAΦNE the ϕ is produced with a small boost in the horizontal plane the kaons have different values for γ_K as a function of the azimuthal angle. The ratio of the decay intensities $I_{\pm}(\Delta t)$ distribution for events in which the K_+ propagates opposite or along the ϕ momentum will enhance the small asymmetry introduced by the Δa_0 component of the δ parameter. For values of Δa_0 of the order $\mathcal{O}(10^{-18})$ we expect an effect on the $I_{\pm}(\Delta t)$ up to 1%-2% in the region $|\Delta t| < 5\tau_S$ from which we expect to be able to put limits of the order $\mathcal{O}(10^{-17})$ - $\mathcal{O}(10^{-18})$.

The Δa_{μ} parameters can be all simultaneously measured by performing a proper sidereal time dependent analysis of asymmetries of $I_{\pm}(\Delta t)$, Eq. (2) and Eq. (5). An accuracy $\mathcal{O}(10^{-18})$ GeV could be reached with the analysis of the full KLOE data sample.

3. Conclusions and future plans

All four Δa_{μ} parameters of the SME can be independently measured at KLOE, completing results obtained by fixed beam experiments.

The continuation of the KLOE physics program with KLOE-2⁹ at an improved DAΦNE machine is currently starting. The data taking campaign

will be organized in two different steps. During the first one we will double the statistics already taken by KLOE with a new beam interaction scheme¹⁰ and with the inclusion of two pairs of electron-positron taggers¹¹ for the study of the gamma-gamma physics. The second phase aims to an integrated luminosity of $\sim 25 \text{ fb}^{-1}$ including several upgrades for the KLOE detector:

- a pair of crystal calorimeters (CCALT¹²) near the interaction region to improve the angular acceptance for low- θ particles;
- a pair of tile calorimeters (QCALT¹³) covering the quadrupoles along the beam pipe made of tungsten foil and singly read-out scintillator tiles to improve the angular coverage for particles coming from the active volume of the DC;
- a small and light tracker (IT¹⁴) made of four planes of cylindrical GEM to improve the resolution of the vertex reconstruction around the interaction point and to increase the low- θ charged particles acceptance.

One of the main physics issue of KLOE-2 is the search for CPT violation effects; limits on several parameters are expected to be improved by about one order of magnitude.

References

1. M. Adinolfi *et al.*, KLOE Collaboration, Nucl. Inst. Meth. A **488** (2002) 51.
2. M. Adinolfi *et al.*, KLOE Collaboration Nucl. Inst. Meth. A **482** (2002) 363.
3. V.A. Kostelecký, Phys. Rev. Lett. **80**, 1818 (1998).
4. V.A. Kostelecký, Phys. Rev. D **61**, 016002 (1999).
5. V.A. Kostelecký, Phys. Rev. D **64**, 076001 (2001).
6. A. Di Domenico, in V.A. Kostelecký, ed., *Fourth Meeting on CPT and Lorentz Symmetry*, World Scientific, Singapore, 2008; A. Di Domenico, in A. Di Domenico ed., *Handbook on neutral kaon interferometry at a Φ -factory*, Frascati Physics Series, Vol. 43.
7. C. D. Buchanan *et al.*, Phys. Rev. D **45**, 4088 (1992).
8. A. Di Domenico, J. Phys. Conf. Ser. **171**, 012008 (2009).
9. G. Amelino-Camelia *et al.*, arXiv:1003.3868.
10. C. Milardi *et al.*, DAFNE Collaboration, arXiv:1006.1487.
11. D. Babusci *et al.*, Nucl. Instrum. Meth. A **617**, 81 (2010).
12. F. Happacher *et al.*, Nucl. Phys. Proc. Suppl. **197**, 215 (2009).
13. M. Cordelli *et al.*, Nucl. Instrum. Meth. A **617**, 105 (2010).
14. F. Archilli *et al.*, KLOE-2 Collaboration, arXiv:1002.2572.